

- ✓ High potency methionine injection
- ✓ An essential amino acid
- ✓ Methionine – converts to cystine immediate source muscle energy
- ✓ Protects the liver – a liver tonic
- ✓ Acidifies urine


DL-Acetyl-Methionine

INJECTABLE AMINO ACID

DL-Acetyl-Methionine 200 mg/mL

Methionine is a lipotropic and is liver protecting. It is an essential amino acid containing a sulfa atom which protects the liver against necrosis. It also protects and treats steatosis (retention of lipids in the liver).

Methionine is a precursor of the amino acid cystine; which is an important source of muscle energy. It acidifies urine.




Protects and de-toxifies the liver

Methionine protects the liver against injury, detoxifies and aids repair of liver damage. It is a lipotropic agent, which plays a role in healthy liver function.

Methionine assists in the breakdown of fats in the liver (fatty liver disease – cattle).

It has antioxidant properties, which protects the liver, lymphatic system and other tissues in the body against free-radical formation and damage. This novel, indirect antioxidant action might be relevant for prevention of inflammation and oxidative stress.



ausrichter

DL-Acetyl-Methionine

Injectable Amino Acid Supplement

PRODUCT INFORMATION

Composition

Acetyl-dl-methionine 200 mg/mL / 50 mL.

Actions

Hepto-protective, detoxification of the liver, liver insufficiency supportive treatment of liver disease (protection of the formation of free-radicals in tissues).

Indications

For use as supportive therapy in the treatment of liver disease in horses cattle, sheep, goats, pigs, dogs and cats.

Complementary treatment of –

- ✓ Hepatitis
- ✓ Hepatic steatosis
- ✓ Metabolic disorders
- ✓ Acetonemia
- ✓ Hypocalcemia
- ✓ Tetany
- ✓ For acidification of urine

Directions for use

By slow intramuscular (IM) or IV injection.

Dosage

10 mL per 100 kg bodyweight;
Dogs 10 mL per 20 kg;
Cats 1 mL per kg bodyweight.

Repeat after one week or as directed by a veterinarian.

The intramuscular IM injection is sometimes slightly painful. To avoid this the required dose may be diluted in twice the volume of the injection dose in sterile water.

Withholding period

NIL Racing. It is preferable to suspend administration 48 hours before racing.

Export slaughter interval: NIL.

Presentation

Injection 50 mL.

Storage

Store below 25°C.

APVMA No. 36730 / 50 mL / 0108

ausrichter

Ausrichter Pty Ltd

ABN 79 000 908 529

2/21 Chester Street, Camperdown NSW 2050

Telephone: (02) 9517 1166

Fax: (02) 9516 5810

Email: info@ausrichter.com

www.ausrichter.com

Animal Health Products